

=> d his

(FILE 'HOME' ENTERED AT 16:30:03 ON 21 NOV 2007)


FILE 'CAPLUS, MEDLINE' ENTERED AT 16:33:52 ON 21 NOV 2007

L1	138 S	PYRIDOXINE (P) ?GLUCOS?	(P) SYNTH?
L2	0 S	L1 AND LEAVING GROUP?	
L3	0 S	L1 AND HALOGEN?	
L4	0 S	L1 AND HALIDE?	
L5	28 S	L1 AND ?THIO?	
L6	110 S	L1 NOT L5	
L7	18 S	L6 AND ?GLUCOSIDE?	

=> d 11


L1 HAS NO ANSWERS

L1 STR




Structure attributes must be viewed using STN Express query preparation.

=> d 11  
L1 HAS NO ANSWERS  
L1 STR



Structure attributes must be viewed using STN Express query preparation.



chain nodes :

7 8 9 10 11 12 13

ring nodes :

1 2 3 4 5 6 14 15 16 17 18 19 20 21 22 23 24 25

chain bonds :

2-9 3-10 4-11 5-7 6-13 7-8 10-18 11-12

ring bonds :

1-2 1-6 2-3 3-4 4-5 5-6 14-15 14-19 15-16 16-17 17-18 18-19 20-21 20-25 21-22 22-23 23-24 24-25

exact/norm bonds :

3-10 10-18 14-15 14-19 15-16 16-17 17-18 18-19 20-21 20-25 21-22 22-23 23-24 24-25

exact bonds :

2-9 4-11 5-7 6-13 7-8 11-12

normalized bonds :

1-2 1-6 2-3 3-4 4-5 5-6

Match level :


1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:CLASS8:CLASS9:CLASS10:CLASS11:CLASS12:CLASS13:CLASS14:Atom 15:Atom 16:Atom 17:Atom 18:Atom 19:Atom 20:Atom 21:Atom 22:Atom 23:Atom 24:Atom 25:Atom

fragments assigned product role:

containing 1

fragments assigned reactant/reagent role:





chain nodes :

7 8 9 10 11 30 31 32 33 34

ring nodes :

1 2 3 4 5 6 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29

chain bonds :

2-9 3-10 5-7 6-11 7-8 10-16 25-32 26-33 28-30 29-34 30-31

ring bonds :

1-2 1-6 2-3 3-4 4-5 5-6 12-13 12-17 13-14 14-15 15-16 16-17 18-19 18-23 19-20 20-21 21-22 22-23 24-25 24-29 25-26 26-27 27-28 28-29

exact/norm bonds :

3-10 10-16 12-13 12-17 13-14 14-15 15-16 16-17 18-19 18-23 19-20 20-21 21-22 22-23 26-33

exact bonds :

2-9 5-7 6-11 7-8 25-32 28-30 29-34 30-31

normalized bonds :

1-2 1-6 2-3 3-4 4-5 5-6 24-25 24-29 25-26 26-27 27-28 28-29

Match level :



1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:CLASS8:CLASS9:CLASS10:CLASS11:CLASS12:Atom  
13:Atom 14:Atom 15:Atom 16:Atom 17:Atom 18:Atom 19:Atom 20:Atom 21:Atom 22:Atom 23:Atom  
24:Atom 25:Atom 26:Atom 27:Atom 28:Atom 29:Atom 30:CLASS31:CLASS32:CLASS33:CLASS34:CLASS

fragments assigned product role:

containing 1

fragments assigned reactant/reagent role:

containing 24



chain nodes :

7 8 9 10 11

ring nodes :

1 2 3 4 5 6 12 13 14 15 16 17 18 19 20 21 22 23

chain bonds :

2-9 3-10 5-7 6-11 7-8 10-16

ring bonds :

1-2 1-6 2-3 3-4 4-5 5-6 12-13 12-17 13-14 14-15 15-16 16-17 18-19 18-23 19-20 20-21 21-22 22-23

exact/norm bonds :

3-10 10-16 12-13 12-17 13-14 14-15 15-16 16-17 18-19 18-23 19-20 20-21 21-22 22-23

exact bonds :

2-9 5-7 6-11 7-8

normalized bonds :

1-2 1-6 2-3 3-4 4-5 5-6

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:CLASS8:CLASS9:CLASS10:CLASS11:CLASS12:Atom  
13:Atom 14:Atom 15:Atom 16:Atom 17:Atom 18:Atom 19:Atom 20:Atom 21:Atom 22:Atom 23:Atom

fragments assigned product role:

containing 1

fragments assigned reactant/reagent role:

containing 18



L4 ANSWER 1 OF 6 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2005:324175 CAPLUS

DOCUMENT NUMBER: 142:397731

TITLE: Stable vitamin B6 derivatives

INVENTOR(S): Sakamoto, Keiji; Wada, Koichi; Ito, Hajime; Take, Nobuhiro; Morimoto, Hiroshi; Maniwa, Fumio; Shimmoto, Yukiko

PATENT ASSIGNEE(S): Daiichi Fine Chemical Co., Ltd., Japan

SOURCE: PCT Int. Appl., 64 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent


LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005033123	A1	20050414	WO 2004-JP14768	20040930
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
CA 2544574	A1	20050414	CA 2004-2544574	20040930
EP 1679316	A1	20060712	EP 2004-773642	20040930
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK			
CN 1863811	A	20061115	CN 2004-80028719	20040930
IN 2006CN01487	A	20070629	IN 2006-CN1487	20060501
US 2007148108	A1	20070628	US 2007-573973	20070222
PRIORITY APPLN. INFO.:			JP 2003-342918	A 20031001
			JP 2004-155624	A 20040526
			WO 2004-JP14768	W 20040930

GI



I

AB Disclosed are pyridoxine derivs. (I) (wherein R1 represents a glycosyl group, a phosphoric acid group, or a cyclic phosphoric acid group bonded with R2; R2 represents -CH2OH, -CHO, -CH2NH2, -CH2-amino acid residue or -CH2-OP(=O)(OH)2; and R3 represents a hydrogen atom or -PO3H2) or a salt thereof. Also disclosed is a composition for cosmetics, drugs, foods and/or animal feed which contains such a compound or a salt thereof. Pyridoxine 3-β-D-glucoside was prepared and tested for photostability and heat stability. Pyridoxine 3-β-D-glucoside was used in formulating lotions, shampoos, eye drops, beverages, etc.

IT 72551-78-1P 849790-03-OP 849790-10-9P


849790-11-0P

RL: COS (Cosmetic use); FFD (Food or feed use); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(preparation of stable vitamin B6 derivs. for use in cosmetic and food and pharmaceutical compns.)

RN 72551-78-1 CAPLUS

CN  $\beta$ -D-Glucopyranoside, 4,5-bis(hydroxymethyl)-2-methyl-3-pyridinyl  
(9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 849790-03-0 CAPLUS

CN  $\beta$ -D-Galactopyranoside, 4,5-bis(hydroxymethyl)-2-methyl-3-pyridinyl  
(9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 849790-10-9 CAPLUS

CN  $\beta$ -D-Glucopyranoside, 4,5-bis(hydroxymethyl)-2-methyl-3-pyridinyl,  
hydrochloride (9CI) (CA INDEX NAME)


Absolute stereochemistry.



● HCl

RN 849790-11-0 CAPLUS  
CN  $\alpha$ -D-Glucopyranoside, 4,5-bis(hydroxymethyl)-2-methyl-3-pyridinyl  
(9CI) (CA INDEX NAME)


Absolute stereochemistry.



IT 849790-04-1P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
(Reactant or reagent)  
(preparation of stable vitamin B6 derivs. for use in cosmetic and food and  
pharmaceutical compns.)

RN 849790-04-1 CAPLUS  
CN  $\beta$ -D-Galactopyranoside, 4,5-bis(hydroxymethyl)-2-methyl-3-pyridinyl,  
2,3,4,6-tetraacetate (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 2 OF 6 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1992:193047 CAPLUS

DOCUMENT NUMBER: 116:193047

TITLE: Pyridoxine-5'- $\beta$ -D-glucoside affects the metabolic  
utilization of pyridoxine in rats

AUTHOR(S): Gilbert, Joyce A.; Gregory, Jesse F., III  
CORPORATE SOURCE: Food Sci. Hum. Nutr. Dep., Univ. Florida, Gainesville,  
FL, 32611-0163, USA

SOURCE: Journal of Nutrition (1992), 122(4), 1029-35  
CODEN: JONUAI; ISSN: 0022-3166

DOCUMENT TYPE: Journal  
LANGUAGE: English

AB A major form of vitamin B-6 in plant-derived foods is pyridoxine-5'- $\beta$ -D-glucoside. Previous studies have shown that pyridoxine-5'- $\beta$ -D-glucoside is poorly available as a source of vitamin B-6 in rats and is partially utilized in humans. This research was conducted to determine whether unlabeled pyridoxine-5'- $\beta$ -D-glucoside affects the metabolic utilization of simultaneously administered isotopically labeled pyridoxine in rats. Three groups of rats were administered a single oral dose of 0,

36, or 72 nmol of unlabeled pyridoxine-5'- $\beta$ -D-glucoside along with 166.5 MBq (240 nmol) of [14C]pyridoxine. Twenty-four hours after administration of the dose the rats were killed, and the isotopic distribution of vitamin B-6 metabolites in liver and urine was determined. Urinary 14C and hepatic 14C-labeled pyridoxine phosphate and pyridoxal phosphate were directly related to pyridoxine-5'- $\beta$ -D-glucoside dose. Hepatic 14C, 14C-labeled pyridoxal, pyridoxine and pyridoxamine, and the concentration of urinary [14C]4-pyridoxic acid, relative to total urinary 14C, were inversely proportional to the dose of pyridoxine-5'- $\beta$ -D-glucoside. These results provide evidence that pyridoxine-5'- $\beta$ -D-glucoside quant. alters the metabolism and in vivo retention of [14C]pyridoxine and that pyridoxine-5'- $\beta$ -D-glucoside may retard the utilization of nonglycosylated forms of vitamin B-6.


IT 72551-78-1

RL: BIOL (Biological study)  
(pyridoxine metabolic utilization response to dietary)

RN 72551-78-1 CAPLUS

CN  $\beta$ -D-Glucopyranoside, 4,5-bis(hydroxymethyl)-2-methyl-3-pyridinyl  
(9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 3 OF 6 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1991:19895 CAPLUS

DOCUMENT NUMBER: 114:19895

TITLE: Hydrolysis of pyridoxine-5'- $\beta$ -D-glucoside by a broad-specificity  $\beta$ -glucosidase from mammalian tissues

AUTHOR(S): Trumbo, Paula R.; Banks, Melanie A.; Gregory, Jesse F., III

CORPORATE SOURCE: Food Sci. Hum. Nutr. Dep., Univ. Florida, Gainesville, FL, 32611-0163, USA

SOURCE: Proceedings of the Society for Experimental Biology and Medicine (1990), 195(2), 240-6  
CODEN: PSEBAA; ISSN: 0037-9727

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Research was conducted to evaluate the ability of a broad-specificity  $\beta$ -glucosidase in mammalian tissues to catalyze the hydrolytic release of free pyridoxine from pyridoxine-5'- $\beta$ -D-glucoside, a naturally occurring form of vitamin B6 in plant-derived foods. Activity was detected in liver and intestinal mucosa using tritiated pyridoxine glucoside as a substrate. In the rat and guinea pig, enzyme activity was greater in intestine than in liver or kidney while even greater activity was detected in human intestinal tissue. Reaction rates were, however, low in all tissues. Hydrolysis of the synthetic substrate 4-methylumbelliferone-3-O- $\beta$ -D-glucoside was also greatest in intestinal tissue. The characteristics of the enzymic hydrolysis of pyridoxine glucoside to pyridoxine included: (1) most activity in the soluble tissue fraction, (2) a pH optimum of approx. 6.0, and (3) inhibition caused by the addition of Na taurocholate. These characteristics are very similar to

those of the broad-specificity  $\beta$ -glucosidase in mammalian tissues with respect to the hydrolysis of a variety of naturally occurring and synthetic substrates. The apparent  $K_m$  was greater than 2 mM for pyridoxine glucoside hydrolysis by intestinal preps. of each species, which is much greater than expected intestinal concns. derived from dietary sources. In vivo studies have indicated that the intestine is involved in the metabolic utilization of dietary pyridoxine glucoside. The results observed here suggest that an alternate process, possibly involving intestinal microorganisms, may also be involved in the in vivo hydrolysis of pyridoxine glucoside.

IT 72551-78-1


RL: RCT (Reactant); RACT (Reactant or reagent)

(hydrolysis of, by broad-specificity glucosidase from human and mammal tissue)

RN 72551-78-1 CAPLUS

CN  $\beta$ -D-Glucopyranoside, 4,5-bis(hydroxymethyl)-2-methyl-3-pyridinyl  
(9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 4 OF 6 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1990:54135 CAPLUS

DOCUMENT NUMBER: 112:54135

TITLE: Dietary intake of total and glycosylated vitamin B6 and the vitamin B6 nutritional status of unsupplemented lactating women and their infants

AUTHOR(S): Andon, Mark B.; Reynolds, Robert D.; Moser-Veillon, Phyllis B.; Howard, M. Pat

CORPORATE SOURCE: Dep. Hum. Nutr. Food Syst., Univ. Maryland, College Park, MD, USA

SOURCE: American Journal of Clinical Nutrition (1989), 50(5), 1050-8

CODEN: AJCNAC; ISSN: 0002-9165

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The mean dietary intakes of total and glycosylated vitamin B6, determined from anal. of 3-day diet composites collected from lactating women, were 8.63 and 1.33  $\mu$ mol/day, resp. A comparison of linear regression models that either included or excluded dietary glycosylated vitamin B6 content indicates that the intake of glycosylated vitamin B6 had little, if any, effect upon maternal plasma pyridoxal 5'-phosphate concentration and maternal urinary excretion of total vitamin B6 and 4-pyridoxic acid. On the basis of guidelines from the literature for evaluating biochem. indexes of vitamin B6 nutriture, the women appeared to be consuming adequate amts. of the vitamin. The mean breast-milk concns. of total and glycosylated vitamin B6 were 733 and 18 mM, resp. The infant plasma pyridoxal 5'-phosphate concentration was 54 nM and all infants had lengths and wts. appropriate for their age.


IT 72551-78-1

RL: BIOL (Biological study)

(nutritional status of, in lactating women and their infants)

RN 72551-78-1 CAPLUS  
CN  $\beta$ -D-Glucopyranoside, 4,5-bis(hydroxymethyl)-2-methyl-3-pyridinyl  
(9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 5 OF 6 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1988:185548 CAPLUS

DOCUMENT NUMBER: 108:185548

TITLE: Incomplete utilization of pyridoxine- $\beta$ -glucoside  
as vitamin B6 in the rat

AUTHOR(S): Trumbo, Paula R.; Gregory, Jesse F., III; Sartain,  
Doris B.

CORPORATE SOURCE: Food Sci. Hum. Nutr. Dep., Univ. Florida, Gainesville,  
FL, 32611, USA

SOURCE: Journal of Nutrition (1988), 118(2), 170-5

CODEN: JONUAI; ISSN: 0022-3166

DOCUMENT TYPE: Journal

LANGUAGE: English

AB This research was conducted to determine the bioavailability of 5'-O-( $\beta$ -D-glucopyranosyl) pyridoxine (PN-glucoside) during chronic administration in a depletion-repletion bioassay. PN-glucoside was found previously to constitute a major portion of the total vitamin B6 in many foods of plant origin. Following a 14-day depletion period, rats were fed diets containing graded levels of either free pyridoxine (PN) or PN-glucoside for 17-days. Slope ratio anal. of dose-response curves, on the basis of growth and plasma pyridoxal 5-phosphate (PLP) concentration, indicated 10-34% utilization of PN-glucoside relative to the molar response to PN. Erythrocyte aspartate aminotransferase (AspAT) activity and urinary 4-pyridoxic acid concentration were lower and the stimulation of AspAT activity by exogenous PLP was greater for rats fed PN-glucoside than for those fed PN, which indicated reduced vitamin B6 nutriture in response to PN-glucoside. A constant 7-9% of the ingested PN-glucoside was detected in urine in intact form at all dosage levels. These results provide further evidence of noncomplete bioavailability of PN-glucoside and indicate that its extent of utilization is not influenced by its level of dietary intake.

IT 72551-78-1


RL: PROC (Process)

(bioavailability of, as vitamin B6 source)

RN 72551-78-1 CAPLUS

CN  $\beta$ -D-Glucopyranoside, 4,5-bis(hydroxymethyl)-2-methyl-3-pyridinyl  
(9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 6 OF 6 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1980:55172 CAPLUS

DOCUMENT NUMBER: 92:55172

TITLE: A particulate glucosyltransferase catalyzing the formation of 5'-O-( $\beta$ -D-glucopyranosyl)pyridoxine from pyridoxine: the occurrence in the seedlings of *Pisum sativum* L

AUTHOR(S): Tadera, Kenjiro; Nakamura, Mahomi; Yagi, Fumio; Kobayashi, Akira

CORPORATE SOURCE: Fac. Agric., Kagoshima Univ., Kagoshima, 890, Japan

SOURCE: Journal of Nutritional Science and Vitaminology (1979), 25(4), 347-50

CODEN: JNSVA5; ISSN: 0301-4800

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The 20,000-50,000 g particulate fraction obtained from pea seedlings with a protein concentration of 20 mg/mL catalyzed the glucosylation of pyridoxine. The rate of glucosylation was linear with time for  $\geq 40$  min and proportional to the protein concentration at  $\leq 20$  mg/mL. The pH optimum, determined, in several different buffer systems, was between 7.8 and 8.8. Apparent  $K_m$  values were 0.4 and 0.7 mM for pyridoxine and UDP-glucose resp. The 5'-O-( $\beta$ -D-glucopyranosyl)pyridoxine reaction product, purified by Sephadex G-10 gel filtration and by paper chromatog., was confirmed by chemical tests and Rf value detns.

IT 72551-78-1


RL: FORM (Formation, nonpreparative)

(formation of, from pyridoxine, pea particulate glucosyltransferase catalysis of)


RN 72551-78-1 CAPLUS

CN  $\beta$ -D-Glucopyranoside, 4,5-bis(hydroxymethyl)-2-methyl-3-pyridinyl (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RX(2) OF 4



REF: Gazzetta Chimica Italiana, 119(1), 63-4; 1989

ACCESSION NUMBER:	111:36634 CASREACT
TITLE:	Isolation of a new compounds related to 4-methoxypyridoxine from Albizzia lucida
AUTHOR(S):	Orsini, Fulvia; Pelizzoni, Francesca; Pulici, Maurizio; Verotta, Luisella
CORPORATE SOURCE:	Dip. Chim. Org. Ind., CNR, Milano, I-20133, Italy
SOURCE:	Gazzetta Chimica Italiana (1989), 119(1), 63-4
DOCUMENT TYPE:	CODEN: GCITA9; ISSN: 0016-5603
LANGUAGE:	Journal
GI	English




I

AB A new 32-O-glucoside (I) of 3-hydroxy-5-(hydroxymethyl)-4-(methoxymethyl)-2-methylpyridine was isolated from seeds of Albizzia lucida and its structure determined on the basis of hydrolysis and spectral evidence.




RX(2) OF 4



REF: Gazzetta Chimica Italiana, 119(1), 63-4; 1989

ACCESSION NUMBER: 111:36634 CASREACT  
 TITLE: Isolation of a new compounds related to  
 4-methoxypyridoxine from Albizzia lucida  
 Orsini, Fulvia; Pelizzoni, Francesca; Pulici,  
 Maurizio; Verotta, Luisella  
 CORPORATE SOURCE: Dip. Chim. Org. Ind., CNR, Milano, I-20133, Italy  
 SOURCE: Gazzetta Chimica Italiana (1989), 119(1), 63-4  
 CODEN: GCITA9; ISSN: 0016-5603  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 GI



I

AB A new 32-O-glucoside (I) of 3-hydroxy-5-(hydroxymethyl)-4-(methoxymethyl)-  
 2-methylpyridine was isolated from seeds of Albizzia lucida and its  
 structure determined on the basis of hydrolysis and spectral evidence.

L7 ANSWER 9 OF 18 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1978:49161 CAPLUS

DOCUMENT NUMBER: 88:49161

ORIGINAL REFERENCE NO.: 88:7759a,7762a

TITLE: Isolation from rice bran of a bound form of vitamin B6 and its identification as 5'-O-( $\beta$ -D-glucopyranosyl)pyridoxine

AUTHOR(S): Yasumoto, Kyoden; Tsuji, Hideaki; Iwami, Kimikazu; Mitsuda, Hisateru

CORPORATE SOURCE: Fac. Agric., Kyoto Univ., Kyoto, Japan

SOURCE: Agricultural and Biological Chemistry (1977), 41(6), 1061-7

CODEN: ABCHA6; ISSN: 0002-1369

DOCUMENT TYPE: Journal

LANGUAGE: English

AB One of the bound forms of vitamin B6 [8059-24-3] occurring in rice bran was isolated in a faintly yellowish syrup by repeating ion-exchange and paper-partition chromatog. techniques. The behaviors of the isolate on thin-layer and Aminex A-5 column chromatograms were coincident with those of synthetic pyridoxine  $\beta$ -D- glucoside, which was obtained by Koenigs-Knorr condensation of  $\alpha$ 4,3-O-isopropylidene pyridoxine and 2,3,4,6-tetra-O-acetyl- $\alpha$ -D-glucopyranosyl bromide. On acid hydrolysis, the isolate gave pyridoxine and glucose. Glucose bound to the 5-hydroxymethyl group of pyridoxine, because the isolate did not react with 2,6-dichloroquinone chlorimide in the presence of boric acid. An equimolar amount of pyridoxine and D-glucose was produced with an equivalent consumption of the isolate by the action of  $\beta$ - glucosidase. No essential difference between the isolated and synthetic preps. could be detected in UV and NMR spectra. Thus, the chemical structure of the isolate was 5'-O-( $\beta$ -D-glucopyranosyl) pyridoxine [19316-63-3].

L7 ANSWER 10 OF 18 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1977:466953 CAPLUS

DOCUMENT NUMBER: 87:66953

ORIGINAL REFERENCE NO.: 87:10639a,10642a

TITLE: Availability as vitamin B6 and small intestinal absorption of pyridoxine- $\beta$ -D- glucoside in rats

AUTHOR(S): Tsuji, Hideaki; Okada, Jungo; Iwami, Kimikazu; Yasumoto, Kyoden

CORPORATE SOURCE: Fac. Agric., Kyoto Univ., Kyoto, Japan

SOURCE: Bitamin (1977), 51(4), 153-9

CODEN: BTMNA7; ISSN: 0006-386X

DOCUMENT TYPE: Journal

LANGUAGE: Japanese

AB Utilization of a chemical synthesized pyridoxine  $\beta$ -D- glucoside [63245-12-5] by vitamin B6 [8059-24-3]-deficient rats was examined in terms of its effects on the urinary excretion of xanthurenic acid [59-00-7] and on the activation of vitamin b6 enzymes, glutamate-pyruvate transaminase [9000-86-6] and cysteine desulfhydrase [9012-96-8]. Oral administration of pyridoxine- $\beta$ - glucoside (30  $\mu$ g/animal/day) for 12 days has led to a complete restoration of the urinary excretion of xanthurenic acid to a normal level. The levels of the enzymic activities in liver and erythrocytes recovered with statistical significance to those in the pos. control rats administered pyridoxine. The  $\beta$ - glucosidase catalyzing hydrolysis of pyridoxine- $\beta$ - glucoside was found in small intestine at a significant level, and at a somewhat lesser level in liver and blood. It thus appears that pyridoxine- $\beta$ - glucoside in vivo substitutes for vitamin B6 by enzymic conversion to free pyridoxine either

before or after absorption in the small intestine. The postabsorptive conversion is supported by permeation of pyridoxine- $\beta$ -glucoside into the serosal side with everted sacs and by the high effectiveness as vitamin B6 of i.v. injected pyridoxine- $\beta$ -glucoside.

L7 ANSWER 11 OF 18 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1973:107995 CAPLUS

DOCUMENT NUMBER: 78:107995

ORIGINAL REFERENCE NO.: 78:17339a,17342a

TITLE: Transglycosidation to vitamin B6 by microorganisms.  
VI. Formation of pyridoxine  
glucoside-synthesizing enzyme  
( $\alpha$ -glucosidase) of *Micrococcus* species  
number 431

AUTHOR(S): Kawai, Fusako; Horii, Takio; Yamada, Hideaki; Ogata, Koichi

CORPORATE SOURCE: Kyoto Res. Inst. Food. Sci., Kyoto Univ., Kyoto, Japan

SOURCE: Agricultural and Biological Chemistry (1972), 36(13),  
2607-9

CODEN: ABCHA6; ISSN: 0002-1369

DOCUMENT TYPE: Journal

LANGUAGE: English

AB A pyridoxine (I) glucoside-synthesizing enzyme (II) purified from *Micrococcus* species number 431 (and, as reported previously, a member of the  $\alpha$ -glucosidase group, also catalyzing the transfer of the glucosyl residue of sucrose (III), maltose (IV), and O- $\alpha$ -D-glucoside to I, to form I glucoside) was formed in substantial amts., in the presence of either III or IV, in a basal medium in which III was replaced by 1 of 4 other C compds. Although glucose and glycerol accelerated growth, they had no effect on the II formation. The transglucosidase activity on a IV-containing medium was increased .apprx.15-fold compared with a glucose-containing medium. I evidently inhibited the growth and formation of II. The optimal concentration

of IV for the formation of II was 3%, and II formation was markedly affected by the initial pH of the medium, the optimal pH being 6.0; no growth was observed at pH 5.0. The II activity in cell exts. attained a maximum after 40 hr of cultivation, accompanied by consumption of II. On longer cultivation, however, II activity gradually decreased. On the other hand, II activity in culture broth was .apprx.10% of that cell exts. II could be induced by a 2nd culture of *Micrococcus* species number 431 on a IV-peptone medium, suggesting that II was also an inducible enzyme, like  $\alpha$ -glucosidase from other microbial sources.

L7 ANSWER 12 OF 18 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1972:109608 CAPLUS

DOCUMENT NUMBER: 76:109608

ORIGINAL REFERENCE NO.: 76:17677a,17680a

TITLE: Transglycosidation to vitamin B6 by microorganisms.  
V. Enzymic properties of pyridoxine  
glucoside-synthesizing enzyme  
( $\alpha$ -glucosidase) of *Micrococcus* species  
Number 431

AUTHOR(S): Kawadi, Fusako; Yamada, Hideaki; Ogata, Koichi

CORPORATE SOURCE: Dep. Agric. Chem., Kyoto Univ., Kyoto, Japan

SOURCE: Agricultural and Biological Chemistry (1971), 35(11),  
1660-7

CODEN: ABCHA6; ISSN: 0002-1369

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Partially and highly purified preps. of a pyridoxine glucoside-synthesizing enzyme from *Micrococcus* Number 431

were stable at pH 7.0 and 0-30°. Maximum activity was at pH 8.0 and 37°. Sucrose, phenyl- $\alpha$ -D-glucoside, and maltose served as glucosyl donors and of the vitamin B6 compds. tested only pyridoxine served as a glucosyl acceptor. The activity was inhibited by p-chloromercuribenzoate and heavy metal ions and somewhat by moniodoacetate. Addition of 2-mercaptoethanol overcame the inhibition by p-chloromercuribenzoate and moniodoacetate. Thus, the enzyme appears to be a glucoside-invertase and a sulphydryl enzyme. The enzyme was not affected by chelating agents and not activated by metal ions.

L7 ANSWER 13 OF 18 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1971:120435 CAPLUS

DOCUMENT NUMBER: 74:120435

ORIGINAL REFERENCE NO.: 74:19435a,19438a

TITLE: Transglycosidation to vitamin B6 by microorganisms.  
IV. Purification of a bacterial enzyme catalyzing pyridoxine glucoside synthesis

AUTHOR(S): Kawai, Fusako; Yamada, Hideaki; Ogata, Koichi

CORPORATE SOURCE: Dep. Agric. Chem., Kyoto Univ., Kyoto, Japan

SOURCE: Agricultural and Biological Chemistry (1971), 35(2), 184-90

CODEN: ABCHA6; ISSN: 0002-1369

DOCUMENT TYPE: Journal

LANGUAGE: English

AB *Sarcina* and *Micrococcus* have the ability to synthesize pyridoxine glucoside. Both pyridoxine-5- $\alpha$ -D-glucoside and pyridoxine-4- $\alpha$ -D-glucoside are produced. The bacteria were grown on sucrose, K phosphate buffer, NaCl, MgSO<sub>4</sub>, and yeast media at pH 7.0. Reducing sugars were determined by the Somogyi-Nelson method, pyridoxine glucoside was separated by paper chromatog. in a BuOH:HOAc:water (4:1:1) solvent, products detected with a uv lamp and extracted from the paper with 50% EtOH 1:12.5% NaOAc for 90 min at 37°. The product was assayed at 470 nm after reaction with diazotized p-aminoacetophenone. Glucosidase and transglucosidase were assayed in phosphate buffer pH 8.0 containing sucrose,  $\beta$ -mercaptoethanol and a suitable amount of enzyme for 30 min at 30°. Reaction was stopped by heating and the amount of reducing sugar was measured. The transglucosidase system also containing pyridoxine. The enzyme was purified by (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> fractionation, DEAE-Sephadex, hydroxylapatite and Sephadex G-100 chromatog. to about 354-fold purification and was homogeneous by polyacrylamide electrophoresis and ultracentrifugation.

L7 ANSWER 14 OF 18 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1971:9712 CAPLUS

DOCUMENT NUMBER: 74:9712

ORIGINAL REFERENCE NO.: 74:1537a,1540a

TITLE: Vitamin B2-glycosides. XXIV. Formation of pyridoxine glycoside-like compounds by enzyme preparations having activity of forming riboflavine glycosides

AUTHOR(S): Suzuki, Yukio; Uchida, Kei; Miyake, Toshio

CORPORATE SOURCE: Ohara Inst. Agric. Biol., Okayama Univ., Okayama, Japan

SOURCE: Bitamin (1970), 42(3), 187-92

CODEN: BTMNA7; ISSN: 0006-386X

DOCUMENT TYPE: Journal

LANGUAGE: Japanese

GI For diagram(s), see printed CA Issue.

AB Pyridoxine glycoside-like compds. were synthesized by partially purified enzyme preps. from *Leuconostoc mesenteroides*, *Aspergillus niger*, and *Mucor javanicus*, and by pure  $\beta$ -galactosidase

from *Escherichia coli* having activity for forming riboflavine glycosides. The formation of riboflavine- $\alpha$ -glucoside in the growth media of *Sarcina* species containing sucrose and riboflavine (I) was closely related to the glucosidation of pyridoxine (II).

L7 ANSWER 15 OF 18 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1969:524851 CAPLUS

DOCUMENT NUMBER: 71:124851

ORIGINAL REFERENCE NO.: 71:23231a,23234a

TITLE: Transglycosidation to vitamin B6 by microorganisms.  
II. Chemical structure of pyridoxine glucoside

AUTHOR(S): Ogata, Koichi; Uchida, Yoshihiro; Kurihara, Norio; Tani, Yoshiki; Tochikura, Tatsurokuro

CORPORATE SOURCE: Kyoto Univ., Kyoto, Japan

SOURCE: Journal of Vitaminology (1969), 15(2), 160-6

CODEN: JVITA5; ISSN: 0022-5398

DOCUMENT TYPE: Journal

LANGUAGE: English

GI For diagram(s), see printed CA Issue.

AB Pyridoxine G (I) was shown to not be a  $\beta$ -D-glucoside by the failure of a  $\beta$ -glucosidase prepared from *Aspergillus niger* (CA 45: 3445d) to hydrolyze it, and is probably an  $\alpha$ -D-glucoside, since it underwent some hydrolysis by an  $\alpha$ -D-glucosidase prepared from brewer's yeast (S. Chiba, et al., 1962). I was dehydrogenated by a pyridoxine dehydrogenase from yeast (CA 55: 7493a) to a pyridoxal, which formed a pyridoxal D-glucoside semicarbazone, which was possible if the main component were pyridoxine 5'- $\alpha$ -D-glucoside (II). I was chromatographed on Dowex.1 + 2 (borate form) into fraction 1, pyridoxine 4'- $\alpha$ -D-glucoside (III), and fraction 2. Acetylation of I with Ac2O and pyridine and treatment with dry HCl gave pyridoxine 4'- $\alpha$ -D-glucoside hexaacetate-HCl, m. 154-7°, and pyridoxine 5'- $\alpha$ -D-glucoside hexaacetate-HCl (IV). A mixture of 1.04 g. isopropylidene pyridoxine (V), 11.0 g. Ag2CO3, 23.0 g. Drierite, and 120 ml. dry benzene was stirred 5 hrs. in the dark, then stirred with 0.3 g. AgClO4 and 2.0 g. 2,3,4,6-tetra-O-benzyl-D-glucopyranosyl chloride in 40 ml. dry benzene until V could no longer be detected by thin-layer chromatog., worked up, and the benzyl groups were removed from the syrupy product by treatment with 50 ml. 85% EtOH containing 1 g. hydrogenated Pd chloride 48 hrs. to yield isopropylidene pyridoxine 5'- $\alpha$ -D-glucose, which was heated with 80% HOAc in a boiling water bath 2 hrs. to give II; the  $\beta$ -D-anomer was removed by exhaustive hydrolysis with the  $\beta$ -D-glucosidase from *Aspergillus niger*. Acetylation of synthetic II gave IV as a colorless syrup. N.M.R. and uv spectral curves were shown.

L7 ANSWER 16 OF 18 MEDLINE on STN

ACCESSION NUMBER: 2003578696 MEDLINE

DOCUMENT NUMBER: PubMed ID: 14660349

TITLE: Use of borate to control the 5'-position-selective microbial glucosylation of pyridoxine.

AUTHOR: Wada Koichi; Asano Yasuhisa

CORPORATE SOURCE: Biotechnology Research Center, Toyama Prefectural University, Kosugi, Toyama 939-0398, Japan..

k-wada@daiichi-fc.jp

SOURCE: Applied and environmental microbiology, (2003 Dec) Vol. 69, No. 12, pp. 7058-62.

Journal code: 7605801. ISSN: 0099-2240.

PUB. COUNTRY: United States

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English

FILE SEGMENT: Priority Journals

ENTRY MONTH: 200404  
ENTRY DATE: Entered STN: 16 Dec 2003  
Last Updated on STN: 17 Apr 2004  
Entered Medline: 16 Apr 2004

AB Nearly 100% 5'-position selectivity of transglucosylation from maltodextrin to pyridoxine (PN) by cells of *Verticillium dahliae* TPU 4900 was observed when the reaction was carried out with borate. The same effect of borate was observed not only during synthesis of pyridoxine 5'-alpha-D-glucoside by partially purified enzyme of this strain but also during synthesis of this compound by other microorganisms and with other enzymes (alpha-glucosidase and cyclomaltodextrin glucanotransferase). The effect was thought to be caused by the formation of a borate complex with 3- and 4'-position hydroxyl groups of PN. A decrease in the formation of pyridoxine 5'-alpha-D-glucoside was observed in the reaction with borate, but this decrease was overcome by optimizing the pH and increasing the amount of cells in the reaction mixture.

L7 ANSWER 17 OF 18 MEDLINE on STN  
ACCESSION NUMBER: 2003203314 MEDLINE  
DOCUMENT NUMBER: PubMed ID: 12723597  
TITLE: Improvement in 5'-position-selective glucosylation of pyridoxine by *Verticillium dahliae* TPU 4900.  
AUTHOR: Wada Koichi; Asano Yasuhisa  
CORPORATE SOURCE: Biotechnology Research Center, Toyama Prefectural University, 5180 Kurokawa, Kosugi, Toyama 939-0398, Japan..  
SOURCE: k-wada@daiichi-fcj.co.jp  
Bioscience, biotechnology, and biochemistry, (2003 Mar)  
Vol. 67, No. 3, pp. 508-16.  
Journal code: 9205717. ISSN: 0916-8451.  
PUB. COUNTRY: Japan  
DOCUMENT TYPE: (COMPARATIVE STUDY)  
Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 200309  
ENTRY DATE: Entered STN: 2 May 2003  
Last Updated on STN: 3 Sep 2003  
Entered Medline: 2 Sep 2003

AB Optimization of culture and reaction conditions for 5'-position-selective transglucosylation to pyridoxine by *Verticillium dahliae* TPU 4900 was investigated. *V. dahliae* TPU 4900 had high transglucosylation activity when grown with soluble starch as a carbon source and organic nitrogens such as Esusan meat as a nitrogen source at 15-20 degrees C. Both the yield of pyridoxine 5'-alpha-D-glucoside (PN-5'-alpha-G) and the 5'-position-selectivity reached a maximum when an intact-cell reaction was done at 50-60 degrees C and pH 7 with additions of dextrin. The transglucosylation activity in culture broth was 71 times with the optimization of culture conditions that under the conditions used for screening. The productivity of PN-5'-alpha-G synthesis was 6.9 times that under the initial conditions when the reaction conditions of intact cells were optimized. From 1000 mM (206 g/L) pyridoxine hydrochloride, PN-5'-alpha-G was synthesized to the concentration of 300 mM (98.4 g/L as PN-5'-alpha-G) with 5'-selectivity of 85% in 53 h by intact cells of *V. dahliae* TPU 4900.

L7 ANSWER 18 OF 18 MEDLINE on STN  
ACCESSION NUMBER: 91046154 MEDLINE  
DOCUMENT NUMBER: PubMed ID: 2122467  
TITLE: Hydrolysis of pyridoxine-5'-beta-D-glucoside by a broad-specificity beta-glucosidase from mammalian tissues.  
AUTHOR: Trumbo P R; Banks M A; Gregory J F 3rd  
CORPORATE SOURCE: Food Science and Human Nutrition Department, University of

CONTRACT NUMBER: Florida, Gainesville 32611-0163.  
DK37481 (NIDDK)  
F32-DK08179 (NIDDK)  
SOURCE: Proceedings of the Society for Experimental Biology and  
Medicine. Society for Experimental Biology and Medicine  
(New York, N.Y.), (1990 Nov) Vol. 195, No. 2, pp. 240-6.  
Journal code: 7505892. ISSN: 0037-9727.  
PUB. COUNTRY: United States  
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
(RESEARCH SUPPORT, U.S. GOV'T, P.H.S.)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 199012  
ENTRY DATE: Entered STN: 8 Feb 1991  
Last Updated on STN: 3 Feb 1997  
Entered Medline: 4 Dec 1990

- AB Research was conducted to evaluate the ability of a broad-specificity beta-glucosidase in mammalian tissues to catalyze the hydrolytic release of free pyridoxine from pyridoxine-5'-beta-D-glucoside, a naturally occurring form of vitamin B6 in plant-derived foods. Activity was detected in liver and intestinal mucosa using tritiated pyridoxine glucoside as a substrate. In the rat and guinea pig, enzyme activity was greater in intestine than in liver or kidney while even greater activity was detected in human intestinal tissue. Reaction rates were, however, low in all tissues. Hydrolysis of the synthetic substrate 4-methylumbelliferyl-beta-D-glucoside was also greatest in intestinal tissue. The characteristics of the enzymatic hydrolysis of pyridoxine glucoside to pyridoxine included: (i) most activity in the soluble tissue fraction, (ii) a pH optimum of approximately 6.0, and (iii) inhibition caused by the addition of sodium taurocholate. These characteristics are very similar to those of the broad-specificity beta-glucosidase in mammalian tissues with respect to the hydrolysis of a variety of naturally occurring and synthetic substrates. The apparent Km was greater than 2 mM for pyridoxine glucoside hydrolysis by intestinal preparations of each species, which is much greater than expected intestinal concentrations derived from dietary sources. In vivo studies have indicated that the intestine is involved in the metabolic utilization of dietary pyridoxine glucoside. The results observed here suggest that an alternate process, possibly involving intestinal microorganisms, may also be involved in the in vivo hydrolysis of pyridoxine glucoside.

L7 ANSWER 4 OF 18 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1996:604932 CAPLUS

DOCUMENT NUMBER: 125:301365

TITLE: Preparation of two pyridoxine- $\alpha$ -glucosides by  $\alpha$ -glucosidase from *Mucor javanicus*

AUTHOR(S): Suzuki, Yukio; Doi, Yusuke; Uchida, Kei; Tsuge, Haruhito

CORPORATE SOURCE: Res. Inst. Bioresour., Okayama Univ., Kurashiki, 710, Japan

SOURCE: Oyo Toshitsu Kagaku (1996), 43(3), 369-372

CODEN: OTKAE3; ISSN: 1340-3494

PUBLISHER: Nippon Oyo Toshitsu Kagakkai

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Two glucosylated compds. of pyridoxine were synthesized in a considerable yield from dextrin and pyridoxine by *Mucor javanicus*  $\alpha$ -glucosidase. The ratio of the products was 1:1. The structures of the products were identified as 5'-O-( $\alpha$ -glucopyranosyl) pyridoxine and 4'-O-( $\alpha$ -glucopyranosyl) pyridoxine by elementary analyses, UV, <sup>1</sup>H- and <sup>13</sup>C-NMR spectra, hydrolysis by  $\alpha$ - and  $\beta$ -glucosidases, migration on paper electrophoresis, and Gibbs reaction in the presence and absence of boric acid.



L7 ANSWER 1 OF 18 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2004:1125170 CAPLUS

DOCUMENT NUMBER: 142:70772

TITLE: Regioselective glucosylation of pyridoxine by microbial glycosyltransferase for pyridoxine 5'- $\alpha$ - glucoside synthesis

INVENTOR(S): Wada, Koichi; Sakamoto, Keiji; Asano, Yasuhisa

PATENT ASSIGNEE(S): Daiichi Fine Chemical Co., Ltd., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 13 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent



LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2004357591	A	20041224	JP 2003-160235	20030605
PRIORITY APPLN. INFO.: OTHER SOURCE(S):	MARPAT 142:70772		JP 2003-160235	20030605

GI



AB A glycosyltransferase capable of catalyzing the conversion of pyridoxine compds. (I) (R = hydrogen atom, lower alkyl group, lower hydroxyalkyl group, carboxyl group, or aldehyde group) to compound (II), derived from microorganisms, and use in enzymic synthesis of pyridoxine 5'- $\alpha$ - glucoside, are disclosed. Pyridoxine-5'- $\alpha$ - glucoside is manufactured by fermentation with microorganism in the presence of boric acid or salt. The pyridoxine is selectively glycosidated at the 5'-position and the byproduct pyridoxine-4'-glucoside is insignificant. II has better photostability, and does not have sour and bitter tastes. It can be absorbed and easily converted to pyridoxal phosphate. Microorganisms from culture collections and isolates from nature were screened for the ability to catalyze the regioselective glucosylation of pyridoxine (PN) to produce pyridoxine 5'- $\alpha$ -D- glucoside (PN-5'- $\alpha$ -G) or pyridoxine 4'- $\alpha$ -D- glucoside (PN-4'- $\alpha$ -G). Transglucosylation activity specific to 5'-position of PN was found in fungi belonging to genera such as *Coriolus* and *Verticillium*, and activity at the 4'-position of PN was found in bacteria belonging to genera such as *Bacillus* and *Serratia*. From 100 mM PN, intact cells of

Verticillium dahliae TPU 4900 produced 42 mM (13.9 mg/mL) PN-5'- $\alpha$ -G after 70 h of reaction. Optimization of culture and reaction conditions for 5'-position-selective transglucosylation to pyridoxine by Verticillium dahliae TPU 4900 was investigated. V. dahliae TPU 4900 had high transglucosylation activity when grown with soluble starch as a carbon source and organic nitrogens such as Esusan

meat

as a nitrogen source at 15-20°C. Both the yield of pyridoxine 5'- $\alpha$ -D-glucoside (PN-5'- $\alpha$ -G) and the 5'-position-selectivity reached a maximum when an intact-cell reaction was done at 50-60°C and pH 7 with addns. of dextrin. The transglucosylation activity in culture broth was 71 times with the optimization of culture conditions that under the conditions used for screening. The productivity of PN-5'- $\alpha$ -G synthesis was 6.9 times that under the initial conditions when the reaction conditions of intact cells were optimized. From 1000 mM (206 g/L) pyridoxine hydrochloride, PN-5'- $\alpha$ -G was synthesized to the concentration of 300 mM (98.4 g/L as PN-5'- $\alpha$ -G) with 5'-selectivity of 85% in 53 h by intact cells of V. dahliae TPU 4900.

L7 ANSWER 2 OF 18 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2003:994589 CAPLUS

DOCUMENT NUMBER: 140:180185

TITLE: Use of borate to control the 5'-position-selective microbial glucosylation of pyridoxine

AUTHOR(S): Wada, Koichi; Asano, Yasuhisa  
CORPORATE SOURCE: Biotechnology Research Center, Toyama Prefectural

SOURCE: University, Toyama, 939-0398, Japan  
Applied and Environmental Microbiology (2003), 69(12), 7058-7062

PUBLISHER: CODEN: AEMIDF; ISSN: 0099-2240  
American Society for Microbiology

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 140:180185

AB Nearly 100% 5'-position selectivity of transglucosylation from maltodextrin to pyridoxine (PN) by cells of Verticillium dahliae TPU 4900 was observed when the reaction was carried out with borate. The same effect of borate was observed not only during synthesis of pyridoxine 5'- $\alpha$ -D-glucoside by partially purified enzyme of this strain but also during synthesis of this compound by other microorganisms and with other enzymes ( $\alpha$ -glucosidase and cyclomaltodextrin glucanotransferase). The effect was thought to be caused by the formation of a borate complex with 3- and 4'-position hydroxyl groups of PN. A decrease in the formation of pyridoxine 5'- $\alpha$ -D-glucoside was observed in the reaction with borate, but this decrease was overcome by optimizing the pH and increasing the amount of cells in the reaction mixture

REFERENCE COUNT: 29 THERE ARE 29 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT.

L7 ANSWER 3 OF 18 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2003:296977 CAPLUS

DOCUMENT NUMBER: 139:35135

TITLE: Improvement in 5'-position-selective glucosylation of pyridoxine by Verticillium dahliae TPU 4900

AUTHOR(S): Wada, Koichi; Asano, Yasuhisa  
CORPORATE SOURCE: Biotechnology Research Center, Toyama Prefectural

SOURCE: University, Toyama, 939-0398, Japan  
Bioscience, Biotechnology, and Biochemistry (2003), 67(3), 508-516

PUBLISHER: CODEN: BBBIEJ; ISSN: 0916-8451  
Japan Society for Bioscience, Biotechnology, and Agrochemistry

DOCUMENT TYPE: Journal  
LANGUAGE: English  
OTHER SOURCE(S): CASREACT 139:35135

AB Optimization of culture and reaction conditions for 5'-position-selective transglucosylation to pyridoxine by *Verticillium dahliae* TPU 4900 was investigated. *V. dahliae* TPU 4900 had high transglucosylation activity when grown with soluble starch as a carbon source and organic nitrogens such as Esusan meat as a nitrogen source at 15-20°C. Both the yield of pyridoxine 5'- $\alpha$ -D-glucoside (PN-5'- $\alpha$ -G) and the 5'-position-selectivity reached a maximum when an intact-cell reaction was done at 50-60°C and pH 7 with addns. of dextrin. The transglucosylation activity in culture broth was 71 times with the optimization of culture conditions that under the conditions used for screening. The productivity of PN-5'- $\alpha$ -G synthesis was 6.9 times that under the initial conditions when the reaction conditions of intact cells were optimized. From 1000 mM (206 g/L) pyridoxine hydrochloride, PN-5'- $\alpha$ -G was synthesized to the concentration of 300 mM (98.4 g/L as PN-5'- $\alpha$ -G) with 5'-selectivity of 85% in 53 h by intact cells of *V. dahliae* TPU 4900.

REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 4 OF 18 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1996:604932 CAPLUS  
DOCUMENT NUMBER: 125:301365  
TITLE: Preparation of two pyridoxine- $\alpha$ -glucosides by  $\alpha$ -glucosidase from *Mucor javanicus*  
AUTHOR(S): Suzuki, Yukio; Doi, Yusuke; Uchida, Kei; Tsuge, Haruhito  
CORPORATE SOURCE: Res. Inst. Bioresour., Okayama Univ., Kurashiki, 710, Japan  
SOURCE: Oyo Toshitsu Kagaku (1996), 43(3), 369-372  
CODEN: OTKAE3; ISSN: 1340-3494  
PUBLISHER: Nippon Oyo Toshitsu Kagakkai  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB Two glucosylated compds. of pyridoxine were synthesized in a considerable yield from dextrin and pyridoxine by *Mucor javanicus*  $\alpha$ -glucosidase. The ratio of the products was 1:1. The structures of the products were identified as 5'-O-( $\alpha$ -glucopyranosyl) pyridoxine and 4'-O-( $\alpha$ -glucopyranosyl) pyridoxine by elementary analyses, UV, 1H- and 13C-NMR spectra, hydrolysis by  $\alpha$ - and  $\beta$ -glucosidases, migration on paper electrophoresis, and Gibbs reaction in the presence and absence of boric acid.

L7 ANSWER 5 OF 18 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1996:485509 CAPLUS  
DOCUMENT NUMBER: 125:245721  
TITLE: Enzymic synthesis of glycosylated and phosphatidylated biologically active compounds  
AUTHOR(S): Suzuki, Yukio; Kim, Young Hoi; Uchida, Kei; Takami, Masaaki  
CORPORATE SOURCE: Res. Inst. Bioresour., Okayama Univ., Kurashiki, 710, Japan  
SOURCE: Oyo Toshitsu Kagaku (1996), 43(2), 273-282  
CODEN: OTKAE3; ISSN: 1340-3494  
PUBLISHER: Nippon Oyo Toshitsu Kagakkai  
DOCUMENT TYPE: Journal; General Review  
LANGUAGE: Japanese

AB A review with 49 refs. The enzymic glycosylation and phosphatidylation of biol. active compds. are described. PhCH<sub>2</sub>OH, 2- or 4-HOC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>OH,

geraniol, and citronellol were glycosylated by incubating with *Aspergillus niger*  $\beta$ -glucosidase in solution containing cellobiose and MeCN, followed by extraction with BuOH, treatment with Amberlite XAD-2 and SiO<sub>2</sub>-gel chromatog. to give  $\beta$ -glucosides of each compound in crystalline state.  $\beta$ -Galactosides of farnesol and geranylgeraniol were obtained in similar manner by treating with *A. oryzae*  $\beta$ -galactosidase in the presence of lactose.  $\beta$ -Glucoside and  $\beta$ -galactoside of tryptophol were also prepared enzymically by incubating tryptophol with resp. enzyme and Ph  $\beta$ -glucoside or o-nitrophenyl  $\beta$ -galactoside. All these  $\beta$ -glycosylated compound were odorless.  $\alpha$ -Type glycosides were prepared by the glucosyl transfer action of bacterial cyclodextrin glucanotransferase (CGTase) from dextrin. Glucosyl transfer was observed not only to CH<sub>2</sub>OH of PhCH<sub>2</sub>OH, and related alcs., riboflavin, pyridoxine, thiamine (B<sub>1</sub>), and BuOH, but also to the OH at the inositol moiety of kasugamycin, at C-4 of glucose moieties of ginsenosides R<sub>c</sub> and R<sub>g</sub>1, at C-3 of fructose, and also to the OH of sec- and tert-Bu alcs. quercetin, vanillin, ethylvanillin, PhOH, pyrocatechol, pyrogallol, gallic acid, and protocatechuic acid, showing broad acceptor specificity of CGTase.  $\alpha$ -Glucosylated compds. of aromatic alcs., vanillin, ethylvanillin, and B<sub>1</sub> were odorless. All glycosylated antioxidants were much more stable than aglycons against oxidation by peroxidase with H<sub>2</sub>O<sub>2</sub>. Enzymic transfer of dipalmitoylphosphatidyl (DDP)-residue from 1,2-dipalmitoyl-3-sn-phosphatidylcholine (DPPC) to the CH<sub>2</sub>OH in vitamins B<sub>1</sub>, B<sub>2</sub>, B<sub>6</sub>, pantothenic acid, B<sub>1</sub> disulfide-related compds., arbutin, kojic acid, genipin, and dihydroxyacetone was studied in order to increase their lipophilic properties. Reactions were carried out by stirring NaOAc buffer containing acceptors and phospholipase D (PLD) from *Streptomyces* with CHCl<sub>3</sub> or EtOAc solution of DPPC at 37°, followed by extraction with CHCl<sub>3</sub> and SiO<sub>2</sub>-gel chromatog. DDP-arbutin and DDP-kojic acid showed the same inhibitory activity to tyrosinase as their parent compds., and DDP-genipin showed 6-52 times stronger cytotoxicity than genipin to HeLa, HEL, and MT-4 cells. DDP-genipin reacted with phenylalanine in organic solvents to give a clear blue solution having a similar color to a natural blue pigment "gardenia blue". Immobilized PLD with Amberlite IRC-50 retained 74% of its initial activity after 10 times-repeated batch reaction for DDP-compound synthesis.

L7 ANSWER 6 OF 18 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1991:19895 CAPLUS

DOCUMENT NUMBER: 114:19895

TITLE: Hydrolysis of pyridoxine-5'- $\beta$ -D- glucoside by a broad-specificity  $\beta$ -glucosidase from mammalian tissues

AUTHOR(S): Trumbo, Paula R.; Banks, Melanie A.; Gregory, Jesse F., III

CORPORATE SOURCE: Food Sci. Hum. Nutr. Dep., Univ. Florida, Gainesville, FL, 32611-0163, USA

SOURCE: Proceedings of the Society for Experimental Biology and Medicine (1990), 195(2), 240-6  
CODEN: PSEBAA; ISSN: 0037-9727

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Research was conducted to evaluate the ability of a broad-specificity  $\beta$ -glucosidase in mammalian tissues to catalyze the hydrolytic release of free pyridoxine from pyridoxine-5'- $\beta$ -D- glucoside, a naturally occurring form of vitamin B<sub>6</sub> in plant-derived foods. Activity was detected in liver and intestinal mucosa using tritiated pyridoxine glucoside as a substrate. In the rat and guinea pig, enzyme activity was greater in intestine than in liver or kidney while even greater activity was detected in human intestinal tissue. Reaction rates were, however, low in all tissues. Hydrolysis of the synthetic substrate 4-methylumbelliferyl- $\beta$ -D- glucoside was also greatest in

intestinal tissue. The characteristics of the enzymic hydrolysis of pyridoxine glucoside to pyridoxine included:

(1) most activity in the soluble tissue fraction, (2) a pH optimum of approx. 6.0, and (3) inhibition caused by the addition of Na taurocholate. These characteristics are very similar to those of the broad-specificity  $\beta$ -glucosidase in mammalian tissues with respect to the hydrolysis of a variety of naturally occurring and synthetic substrates. The apparent  $K_m$  was greater than 2 mM for pyridoxine glucoside hydrolysis by intestinal preps. of each species, which is much greater than expected intestinal concns. derived from dietary sources. In vivo studies have indicated that the intestine is involved in the metabolic utilization of dietary pyridoxine glucoside. The results observed here suggest that an alternate process, possibly involving intestinal microorganisms, may also be involved in the in vivo hydrolysis of pyridoxine glucoside.

L7 ANSWER 7 OF 18 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1988:453491 CAPLUS

DOCUMENT NUMBER: 109:53491

TITLE: Changes in the vitamin B-6 content in potatoes during storage

AUTHOR(S): Addo, Constant; Augustin, Jorg

CORPORATE SOURCE: Dep. Food Sci. Hum. Nutri., Washington State Univ., Pullman, WA, 99164, USA

SOURCE: Journal of Food Science (1988), 53(3), 749-52

CODEN: JFDSA; ISSN: 0022-1147

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The nature of the repeatedly reported increase of vitamin B6 in Russet Burbank potatoes stored for 30 days and 9 mo was investigated. Potatoes contained pyridoxine, pyridoxamine, pyridoxal phosphate, and a pyridoxine glucoside. While pyridoxamine and pyridoxal phosphate concns. remained unchanged, there was a sharp increase in pyridoxine glucose during storage indicating a possible synthesis of vitamin B6 during storage. In general, good agreement existed between the data generated by microbial anal. and those obtained by the HPLC method.

L7 ANSWER 8 OF 18 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1986:477863 CAPLUS

DOCUMENT NUMBER: 105:77863

TITLE: Synthesis of pyridoxine- $\beta$ -glucoside by rice bran  $\beta$ -glucosidase and its in situ absorption in rat small intestine

AUTHOR(S): Iwami, Kimikazu; Yasumoto, Kyoden

CORPORATE SOURCE: Dep. Agric. Chem., Kyoto Prefect. Univ., Kyoto, 606, Japan

SOURCE: Nutrition Research (New York, NY, United States)

(1986), 6(4), 407-14

CODEN: NTRSDC; ISSN: 0271-5317

DOCUMENT TYPE: Journal

LANGUAGE: English

AB A major component of  $\beta$ -glucosidase [9001-22-3] multienzymes was highly purified from rice bran, and by its use, pyridoxine- $\beta$ -glucoside (PIN- $\beta$ -G) was synthesized from p-nitrophenyl- $\beta$ -glucoside [2492-87-7] and pyridoxine [65-23-6]. The synthetic product contained both 4'- [71555-11-8] and 5'-isomers [103584-58-3], which were successfully separated by high-voltage paper electrophoresis. The 4'-isomer was used as a convenient substrate for intestinal absorption of PIN- $\beta$ -G, because it gave a pos. reaction with 2,6-dibromoquinone chlorimide, irresp. of the presence or absence of borate. The absorption

expts. with in situ, isolated rat jejunal loops revealed that the PIN- $\beta$ -G level remaining in the loop did not significantly change within 1 h and that  $\delta$ -gluconolactone, a potent inhibitor of  $\beta$ -glucosidase, did not affect the luminal disappearance of PIN- $\beta$ -G. It thus can be assumed that PIN- $\beta$ -G is absorbed across the intestinal wall by a mechanism of simple diffusion, not by hydrolase-mediated transport.

=> d his

(FILE 'HOME' ENTERED AT 16:30:03 ON 21 NOV 2007)

FILE 'CAPLUS, MEDLINE' ENTERED AT 16:33:52 ON 21 NOV 2007

L1	138 S	PYRIDOXINE (P) ?GLUCOS?	(P) SYNTH?
L2	0 S	L1 AND LEAVING GROUP?	
L3	0 S	L1 AND HALOGEN?	
L4	0 S	L1 AND HALIDE?	
L5	28 S	L1 AND ?THIO?	
L6	110 S	L1 NOT L5	
L7	18 S	L6 AND ?GLUCOSIDE?	